

Transitive actions of locally compact groups on locally contractible spaces

Karl H. Hofmann and Linus Kramer

January 23, 2013

Abstract

Suppose that $X = G/K$ is the quotient of a locally compact group by a closed subgroup. If X is locally contractible and connected, we prove that X is a manifold. If the G -action is faithful, then G is a Lie group.

1 Introduction

In 1974 J. Szenthe stated the following result in [36]. *If a σ -compact locally compact group G , with compact quotient G/G° , acts transitively and faithfully on a locally contractible space X , then X is a manifold and G is a Lie group.* This result, which may be viewed as a solution of Hilbert's 5th problem for transformation groups, has been widely used since then. However, it was discovered in 2011 that Szenthe's proof contains a serious gap. In the present paper we close this gap, proving Szenthe's statement in a different way. Independently and simultaneously, this result was also proved by A.A. George Michael [14]. The last section of our paper contains some more comments on the history of this problem. Our main results are as follows.

Theorem A *Let G be a compact group and let $K \subseteq G$ be a closed subgroup. Suppose that the homogeneous space $X = G/K$ contains a nonempty open subset $V \subseteq X$ which is contractible in X . Then X is a closed manifold. If $N = \bigcap \{gKg^{-1} \mid g \in G\}$ denotes the kernel of the action, then G/N is a compact Lie group acting transitively on X .*

For the case of a locally compact group, we need a stronger topological assumption on the coset space X .

Theorem B *Let G be a locally compact group and let $K \subseteq G$ be a closed subgroup. Suppose that the homogeneous space $X = G/K$ is locally contractible. Then X is a manifold. If X is connected and if $N = \bigcap \{gKg^{-1} \mid g \in G\}$ denotes the kernel of the action, then G/N is a Lie group acting transitively on X .*

In the course of the proof, we need the following extension of Iwasawa's Local Splitting Theorem, which may be interesting in its own right. A local version of this result was proved by Gluškov [15]. This result is also proved in [20] Theorem 4.1 in a different way.

Theorem C *Let G be a locally compact group and let $U \subseteq G$ be a neighborhood of the identity. Then there exist a compact subgroup $N \subseteq U$, a simply connected Lie group L and an open homomorphism $\varphi : N \times L \longrightarrow G$ with discrete kernel, such that $\varphi(n, 1) = n$ for all $n \in N$.*

Conventions and Terminology All maps and group homomorphisms are assumed to be continuous and all spaces and groups are assumed to be Hausdorff unless stated otherwise. Topological countability assumptions will be stated explicitly whenever they are used. The identity component of a topological group G is denoted by G° . This is always a closed normal subgroup. For subgroups $P, Q \subseteq G$ we put

$$\begin{aligned}\text{Cen}_P(Q) &= \{p \in P \mid pq = qp \text{ for all } q \in Q\} \\ \text{Cen}(Q) &= \text{Cen}_Q(Q) \\ \text{Nor}_P(Q) &= \{p \in P \mid pQp^{-1} = Q\} \\ [P, Q] &= \langle [p, q] \mid p \in P \text{ and } q \in Q \rangle\end{aligned}$$

The unit interval is denoted by $[0, 1] = \{t \in \mathbb{R} \mid 0 \leq t \leq 1\}$. For a map $h : X \times [0, 1] \longrightarrow Y$ we write $h_t(x) = h(x, t)$. The projection map of a cartesian product $X \times Y$ is denoted by $\text{pr}_X : (x, y) \longmapsto x$, and similarly $\text{pr}_Y : (x, y) \longmapsto y$.

2 Some homotopy theory of compact groups

We begin by collecting some results which we shall need in the proof of Theorem A. Unless stated otherwise, homotopies are not required to preserve base points. Recall that a map $E \longrightarrow B$ is called a *fibration* if it has the *homotopy lifting property* for every space X . This means that for every commutative diagram

$$\begin{array}{ccc} X \times \{0\} & \xrightarrow{\quad} & E \\ \downarrow & \nearrow \tilde{h} & \downarrow \\ X \times [0, 1] & \xrightarrow{\quad h \quad} & B, \end{array}$$

the dotted lift \tilde{h} exists. The following result is related to the notion of *irreducibility* in [9] p. 394 and in [28].

2.1 Lemma *Let E be a space with the property that every homotopy equivalence $E \xrightarrow{\cong} E$ is surjective. Suppose that $p : E \longrightarrow B$ is a surjective fibration. Then also every homotopy equivalence $B \xrightarrow{\cong} B$ is surjective. If p is homotopic to a map $p' : E \longrightarrow B$, then p' is also surjective.*

Proof. First we note the following. If $\xi : B \longrightarrow B$ is a homotopy equivalence with homotopy inverse η , then $\xi \circ \eta$ is, by definition, homotopic to id_B . In order to show the surjectivity of such a map ξ , it suffices therefore to prove the surjectivity of every map which is homotopic to the identity id_B .

Suppose that $h : B \times [0, 1] \longrightarrow B$ is a map with $h_0 = \text{id}_B$. Then the map $h' : E \times [0, 1] \longrightarrow B$ with $h'_t(x) = h_t(p(x))$ is a homotopy between $p = h'_0$ and $p' = h'_1$. Since E is a fibration, there exists a lift $\tilde{h}' : E \times [0, 1] \longrightarrow E$ of h' , with $\tilde{h}'_0 = \text{id}_E$.

$$\begin{array}{ccc} & & E \\ & \nearrow \tilde{h}' & \downarrow \\ E \times [0, 1] & \xrightarrow{h'} & B \end{array}$$

By our assumptions on E , the map $\tilde{h}'_1 : E \longrightarrow E$ is surjective. It follows that $\tilde{h}'_1(E) = p'(E) = h_1(B) = B$. \square

We now recall several results about the structure of compact groups.

2.2 Theorem (Approximation by compact Lie groups) *Let G be a compact group. Then every neighborhood V of the identity contains a closed normal subgroup $N \trianglelefteq G$ such that G/N is a compact Lie group. The set $\mathcal{N}(G)$ consisting of all closed normal subgroups $N \trianglelefteq G$ such that G/N is a Lie group is a filter basis converging to the identity.*

Proof. See Theorem 9.1 and 2.43 in [21]. \square

2.3 Lemma *Let G be a compact group and $N \trianglelefteq G$ a closed normal subgroup. If N and G/N are Lie groups, then G is a Lie group as well.*

Proof. We show that G has no small subgroups, see [21] Theorem 2.40. Let $W \subseteq G/N$ be a neighborhood of the identity which contains no nontrivial subgroup and let V be its preimage in G under the projection $G \longrightarrow G/N$. Let $U \subseteq G$ be a neighborhood of the identity such that $U \cap N$ does not contain a nontrivial subgroup of N . Then $U \cap V$ contains no nontrivial subgroup of G . \square

2.4 Theorem (Complementation of normal subgroups) *Suppose that G is a compact group and that $N \trianglelefteq G$ is a closed normal subgroup. If G/N is connected, then there exists a closed connected subgroup $M \trianglelefteq G^\circ$ with $G = MN$, and such that $M \cap N$ is totally disconnected.*

Proof. See Theorem 9.77 of [21]. \square

The following lemma about permutations groups is well-known, see [8] Theorem 4.2A. We need, however, a topological version which is somewhat stronger than [32] p. 73.

2.5 Lemma *Suppose that G is a compact group, that K is a closed subgroup and that G acts faithfully on G/K , i.e. that K contains no nontrivial normal subgroup of G . Suppose that $M \subseteq G$ is a closed transitive subgroup, $G = KM$. Then there is an isomorphism of compact groups*

$$\text{Cen}_G(M) \cong \text{Nor}_M(M \cap K) / \text{Cen}_M(M \cap K).$$

Proof. The group $N = \text{Nor}_M(M \cap K)$ acts (from the left) on $X = M/M \cap K$ via

$$n(g(M \cap K)) = g(M \cap K)n^{-1} = gn^{-1}(M \cap K),$$

and the kernel of this action is obviously $Z = \text{Cen}_M(M \cap K)$. The action of N/Z on $M/M \cap K$ is free and it is not difficult to check that N/Z surjects as an abstract group onto the centralizer of M in the symmetric group of X , see [8] Theorem 4.2A or [27] 3.5. The graph

$$\{(g, nZ) \in \text{Cen}_G(M) \times N/Z \mid gK = n^{-1}K\}$$

of the injective homomorphism $\text{Cen}_G(M) \longrightarrow N/Z$ is closed and therefore this map is continuous, see for example [9] XI.2.7. \square

2.6 Theorem *Let M be a compact connected group. If there exists a closed totally disconnected normal subgroup $D \subseteq M$ such that M/D is a Lie group, then there exist simple simply connected compact Lie groups S_1, \dots, S_r and a compact connected finite dimensional abelian group A and a central surjective homomorphism*

$$A \times S_1 \times \dots \times S_r \longrightarrow M$$

with a totally disconnected kernel.

Proof. See [21], Theorem 9.52ff. \square

The following Theorems 2.7 and 2.10 are due to Madison and Mostert [28]. In order to make the paper self-contained, we include proofs.

2.7 Theorem (Madison-Mostert) *Let G be a compact group and let $P, Q \subseteq G$ be closed subgroups, with $P \subseteq Q$. Then the natural map*

$$G/P \longrightarrow G/Q$$

is a fibration.

Proof. Suppose we are given a commutative diagram

$$\begin{array}{ccc} X \times \{0\} & \longrightarrow & G/P \\ \downarrow & \nearrow & \downarrow \\ X \times [0, 1] & \longrightarrow & G/Q. \end{array}$$

We have to show the existence of the dotted map. To this end we consider the poset \mathcal{P} consisting of pairs (φ, N) , where $N \trianglelefteq G$ is a closed normal subgroup and $\varphi : X \times [0, 1] \longrightarrow G/PN$ is a map φ that fits into the commutative diagram

$$\begin{array}{ccccc} X \times \{0\} & \longrightarrow & G/P & \longrightarrow & G/PN \\ \downarrow & & \downarrow & \nearrow & \downarrow \\ X \times [0, 1] & \longrightarrow & G/Q & \longrightarrow & G/QN. \end{array}$$

We put $(\varphi, N) \geq (\psi, M)$ if $N \subseteq M$ and if the diagram

$$\begin{array}{ccccccc} & & G/P & \longrightarrow & G/PN & \longrightarrow & G/PM \\ & & \downarrow & & \downarrow & & \downarrow \\ X \times [0, 1] & \longrightarrow & G/Q & \longrightarrow & G/QN & \longrightarrow & G/QM. \end{array}$$

(Dotted arrows connect $X \times [0, 1]$ to G/P , G/PN , and G/PM .)

commutes. We claim that this partial order is inductive. Suppose that $\mathcal{T} \subseteq \mathcal{P}$ is a linearly ordered subset. Let $L = \bigcap \{N \mid (N, \varphi) \in \mathcal{T}\} \trianglelefteq G$. We have natural maps

$$\begin{array}{ccc} & \prod_{(N, \varphi) \in \mathcal{T}} G/PN & \xleftarrow{\alpha} G/PL \\ & \uparrow \gamma & \nearrow \\ X \times [0, 1] & \xrightarrow{\delta} \prod_{(N, \varphi) \in \mathcal{T}} G/QN & \xleftarrow{\beta} G/QL \\ & \downarrow & \downarrow \end{array}$$

(Dotted arrows connect $X \times [0, 1]$ to G/PL and G/QL .)

Now α and β are injective and hence homeomorphisms onto their respective images. Moreover, the image of γ is contained in the image of α and the image of β is contained in the image of δ . Thus we can fill in the dotted map and obtain an upper bound (L, ψ) of \mathcal{T} . By Zorn's Lemma, \mathcal{P} has maximal elements.

Suppose that $N \trianglelefteq G$ is a compact normal subgroup and that G/N is a Lie group. Therefore the canonical map $G/PN \longrightarrow G/QN$ is a locally trivial fiber bundle, see [37], Theorem 3.58, and hence a fibration, see [9], Theorem 4.2. Thus there exists a map $\varphi : X \times [0, 1] \longrightarrow G/PN$ such that (N, φ) is contained in \mathcal{P} . By Theorem 2.2, there exist arbitrarily small compact normal subgroups $N \trianglelefteq G$ such that G/N is a Lie group. It follows that the maximal elements in \mathcal{P} are of the form $(\{1\}, \varphi)$, and these elements solve the initial lifting problem. \square

As a commentary we mention the following corollary (which will not be used here).

2.8 Corollary *Let G be a compact group and let $P \subseteq Q \subseteq G$ be closed subgroups. Then there is a long exact sequence for the homotopy groups*

$$\longrightarrow \pi_k(Q/P) \longrightarrow \pi_k(G/P) \longrightarrow \pi_k(G/Q) \longrightarrow .$$

\square

Similarly, the Leray-Serre Spectral Sequence may be applied to such a fibration (because every fibration is a Serre fibration). For the next lemma we remark that Čech cohomology and Alexander-Spanier cohomology agree for compact spaces. See also [23] App. II 3.15 and App. III 2.11.

2.9 Lemma *Suppose that a compact totally disconnected group D acts on a compact space X . Then the orbit space map $X \longrightarrow D \backslash X$ induces an injection in Čech cohomology with rational coefficients,*

$$\check{H}^*(X; \mathbb{Q}) \xleftarrow{p} \check{H}^*(D \backslash X; \mathbb{Q}) \longleftarrow 0.$$

Proof. We have by Theorem 2.2 that

$$D = \varprojlim \{D/E \mid E \in \mathcal{N}(D)\}$$

and the groups D/E are finite (in other words, D is a profinite group). It follows that

$$X = \varprojlim \{E \setminus X \mid E \in \mathcal{N}(D)\}.$$

For $E, F \in \mathcal{N}(D)$ with $F \subseteq E$ we have that E/F is finite and thus

$$\check{H}^*(F \setminus X; \mathbb{Q}) \longleftarrow \check{H}^*(E \setminus X; \mathbb{Q})$$

is injective, see [7] Theorem III.7.2. Projective limits of compact spaces commute with Čech cohomology, see [10] X.3. The claim follows now. \square

2.10 Theorem (Madison-Mostert) *Let G be a compact group, let $P \subseteq G$ be a closed subgroup, and let $\xi : G/P \longrightarrow G/P$ be a homotopy equivalence. Then ξ is surjective.*

Proof. As we noted in the proof of Lemma 2.1, we have to show that for every map $h : G/P \times [0, 1] \longrightarrow G/P$ with $h_0 = \text{id}_{G/P}$, the map h_1 is surjective. Let \mathcal{C} denote the class of all pairs (G, P) of compact groups where this holds. Our goal is to show that \mathcal{C} is the class of all compact group pairs.

Claim 1. *If G is a compact connected group, then $(G, 1)$ is in \mathcal{C} . Suppose that this is false. Then there exists a map $h : G \times [0, 1] \longrightarrow G$ with $h_0 = \text{id}_G$, and $h_1 : G \longrightarrow G$ is not surjective. By Theorem 2.2, there exists a closed normal subgroup $N \trianglelefteq G$ such that G/N is a compact connected Lie group, say of dimension r , and an element $g \in G$ such that $gN \cap h_1(G) = \emptyset$. By Theorem 2.4, there exists a compact connected subgroup $M \subseteq G$ such that $G = MN$, and $D = M \cap N$ is totally disconnected. Then $L = M/D \cong G/N$ is a compact connected r -dimensional Lie group. We note that $r > 0$, since otherwise we would have $G = N$. Then have a commutative diagram*

$$\begin{array}{ccccccc} M & \hookrightarrow & G & \xleftarrow{j} & h_1(G) & \xleftarrow{h_1} & G \\ \downarrow & & \downarrow & & \downarrow & & \\ L & \xrightarrow{\cong} & G/N & \longleftarrow & G/N - \{gN\} & & \end{array}$$

Because of the homotopy $h_1 \simeq h_0 = \text{id}_G$, the restriction map $\check{H}^*(G; \mathbb{Q}) \xrightarrow{j^*} \check{H}^*(h_1(G); \mathbb{Q})$ is injective. We have therefore in r -dimensional cohomology a commutative diagram

$$\begin{array}{ccccc} \check{H}^r(M; \mathbb{Q}) & \longleftarrow & \check{H}^r(G; \mathbb{Q}) & \xrightarrow{j^*} & \check{H}^r(h_1(G); \mathbb{Q}) \\ (1) \uparrow & & (2) \uparrow & & \uparrow \\ \check{H}^r(L; \mathbb{Q}) & \xleftarrow{\cong} & \check{H}^r(G/N; \mathbb{Q}) & \longrightarrow & \check{H}^r(G/N - \{gN\}; \mathbb{Q}). \end{array}$$

The map (1) is injective by Lemma 2.9. Thus (2) is also injective. The map j^* is injective by the previous remark. The compact connected Lie group L is \mathbb{Q} -orientable, whence $\check{H}^r(L; \mathbb{Q}) \cong \mathbb{Q}$. On the other hand, $\check{H}^r(G/N - \{gN\}; \mathbb{Q}) = 0$. This is a contradiction.

Claim 2. Suppose that G is a compact group. Then $(G, 1)$ is in \mathcal{C} . Suppose that $a \in G$ is not in the image of $h_1 : G \longrightarrow G$. Then $h'_t(g) = a^{-1}h_t(ag)$ is a map from the identity component G° of G to itself with $h'_0 = \text{id}_{G^\circ}$, and 1 is not in the image of h'_1 , contradicting Claim 1.

Claim 3. If $(G, 1)$ is in \mathcal{C} and if $P \subseteq G$ is closed, then (G, P) is in \mathcal{C} . This follows from Claim 2 and Lemma 2.1, since $G \longrightarrow G/P$ is a fibration by Theorem 2.7. \square

3 The proof of Theorem A

We say that a topological space X is *piecewise contractible* if it satisfies the following condition.

(PC) there is a nonempty open subset $V \subseteq X$ and a continuous map

$$f : V \times [0, 1] \longrightarrow X, \quad (v, t) \longmapsto f_t(v)$$

such that $f_0(v) = v$ for all $v \in V$ and f_1 is a constant map from V into X .

We then call V *contractible in X* , and we say that f *contracts V in X* . We note that the image of f in X is a path-connected set with nonempty interior.

3.1 Lemma *A product space $X \times Y$ is piecewise contractible if and only if X and Y are piecewise contractible.*

Proof. If $V \subseteq X$ and $W \subseteq Y$ are open subsets and if $f : V \times [0, 1] \longrightarrow X$ and $g : W \times [0, 1] \longrightarrow Y$ are as in (PC), then the map $h : V \times W \times [0, 1] \longrightarrow X \times Y$, with $h_t(v, w) = (f_t(v), g_t(w))$ contracts $V \times W$ in $X \times Y$.

Conversely, if $U \subseteq X \times Y$ is a nonempty open subset and if $h : U \times [0, 1] \longrightarrow X \times Y$ contracts U in $X \times Y$, then there is a point $w \in W$ and a nonempty open subset $V \subseteq X$ such that $V \times \{w\} \subseteq U$. Then $f_t(v) = \text{pr}_X(h_t(v, w))$ contracts V in X . \square

We now prove three preparatory lemmas in order to obtain Theorem A.

3.2 Lemma *Let G be a compact group and let $K \subseteq G$ be a closed subgroup. If $X = G/K$ is piecewise contractible, then the subgroup $G^\circ K$ is open in G , the quotient space $G^\circ K/K \cong G^\circ/G^\circ \cap K$ is piecewise contractible, and there is a G° -equivariant homeomorphism*

$$G/K \cong G^\circ K/K \times D$$

for some finite set D .

Proof. The space $G/G^\circ K$ is compact and totally disconnected, see [21] Prop. 10.32 and the following remark. Thus the right action of the compact group $G^\circ K$ on G has a totally disconnected orbit space. Now the result [21] Theorem 10.35 on the Existence of Global Cross

Sections applies and shows that G is homeomorphic to $G^\circ K \times D$ with a totally disconnected compact space D (homeomorphic to $G/G^\circ K$) in such a fashion that the action of $G^\circ K$ is by multiplication on the first factor. In other words, we have a $G^\circ K$ -equivariant homeomorphism $G \cong G^\circ K \times (G/G^\circ K)$, and therefore a G° -equivariant homeomorphism

$$G/K \cong G^\circ K/K \times G/G^\circ K.$$

Now G/K is piecewise contractible. Hence by Lemma 3.1, the totally disconnected compact homogeneous space $D \cong G/G^\circ K$ is piecewise contractible. This implies that each point of D is open, and hence D is finite. So $G^\circ K$ is open in G and $G^\circ K/K$ is open in G/K . Lemma 3.1 implies also that $G^\circ K/K$ is piecewise contractible. \square

3.3 Lemma *Let G be a compact connected group and let $K \subseteq G$ be a closed subgroup. If $X = G/K$ is piecewise contractible, then there exists a closed normal connected subgroup $M \trianglelefteq G$ with $G = KM$ (i.e. M acts transitively on X), and M has a closed central totally disconnected subgroup $D \trianglelefteq M$ such that M/D is a compact connected Lie group.*

Proof. Since $X = G/K$ is homogeneous, we may assume that there is an open neighborhood $V \subseteq G/K$ of the coset $K \in G/K$ and a map $f : V \times [0, 1] \longrightarrow G/K$ as in (PC). By Theorem 2.2 there exists a closed normal subgroup $N \in \mathcal{N}(G)$ with $NK/K \subseteq V$. Let $M \trianglelefteq G$ be a complement of N as in Theorem 2.4. Since $G = MN$, the group N acts transitively on G/KM , and we have by Theorem 2.7 a surjective fibration

$$NK/K \cong N/K \cap N \xrightarrow{p} N/(KM) \cap N \cong G/KM.$$

We define $f' : (N/K \cap N) \times [0, 1] \longrightarrow G/KM$ by $f'_t(n(K \cap N)) = f_t(nK)KM$. Then $f'_0 = p$ and f'_1 is constant. It follows from Lemma 2.1 that $G = KM$. Let $D = M \cap N$. Then $M/D \cong G/N$ is a compact connected Lie group. Since D is totally disconnected and normal in the connected group M , we have that D is central in M . \square

The proof following lemma is partially adapted from [24] Prop. 3.5.

3.4 Lemma *Let G, K, X, M be as in Lemma 3.3. If G acts faithfully on X , then M is a compact Lie group.*

Proof. We put $L = M \cap K$ and we identify X with the quotient space M/L . It follows from Theorem 2.6 that there is a compact connected semisimple Lie group S , a compact connected finite dimensional abelian group A , and a surjective homomorphism $q : A \times S \longrightarrow M$ with a totally disconnected kernel E . Passing to a quotient, we can also assume that E intersects the factors $\{1\} \times S$ and $A \times \{1\}$ trivially. We want to show that A is a Lie group.

Since M acts faithfully and transitively on X , the central subgroup $q(A \times \{1\})$ intersects the stabilizer L trivially. Since X is homogeneous, there is an open neighborhood $V \subseteq X$ of $L \in M/L$ and a map

$$f : V \times [0, 1] \longrightarrow X$$

as in (PC). By Theorem 2.2, there is a closed subgroup $B \subseteq A$ such that $A/B \cong \mathbb{T}^m$ is a compact torus of finite dimension m , and such that $BL/L \subseteq V$. By Lemma 3.3 we may assume

that B is totally disconnected. From the action of $A \times S$ on M/L we have by Theorem 2.7 a fibration

$$A \times S \longrightarrow M/L$$

We also note that B can be identified with its image $BL/L \subseteq M/L$, because $B \cap L = \{1\}$. Then f gives us a map $g : B \times [0, 1] \longrightarrow M/L$, with $g_0(b) = bL$ and g_1 constant. We now lift this to $\tilde{g} : B \times [0, 1] \longrightarrow A \times S$, such that $\tilde{g}_0(b) = b$ for all $b \in B$. We define $h : B \times [0, 1] \longrightarrow A$ by $h_t(b) = \text{pr}_A(\tilde{g}_t(b))$. Now there is a covering map $\varphi : B \times \mathbb{R}^m \longrightarrow A$ with discrete kernel, with $\varphi(b, 0) = b$ for all $b \in B$, see [24] Thm. 13.17 and 13.20. We lift h to a map $\tilde{h} : B \times [0, 1] \longrightarrow B \times \mathbb{R}^m$. The composite

$$B \times [0, 1] \xrightarrow{\tilde{h}} B \times \mathbb{R}^m \xrightarrow{\text{pr}_B} B$$

is a homotopy between id_B and a constant map. Thus B is contractible. On the other hand, B is totally disconnected. Therefore $B = \{1\}$. \square

Now we have collected all ingredients for the proof of the Main Theorem A.

3.5 Theorem *Suppose that G is a compact group and that K is a closed subgroup. Assume also that the action of G on the homogeneous space $X = G/K$ is faithful, i.e. that K contains no nontrivial normal subgroup of G . If X is piecewise contractible, then G is a compact Lie group and X is a closed, but not necessarily connected manifold.*

Proof. By Lemma 3.2 we have a G° -equivariant homeomorphism $G/K \cong G^\circ/G^\circ \cap K \times D$, for some finite set D , and $G^\circ/G^\circ \cap K$ is piecewise contractible. By Lemma 3.4 there exists a closed normal Lie group $M \trianglelefteq G^\circ$ acting transitively on $G^\circ/G^\circ \cap K$. The group G° decomposes as $G^\circ = NM$, where $N \trianglelefteq G$ is a closed normal subgroup that centralizes M . Since G° acts faithfully and transitively on $G^\circ/G^\circ \cap K$, it follows from Lemma 2.5 that N is isomorphic to a closed subgroup of the Lie group $\text{Nor}_M(M \cap K)/\text{Cen}_M(M \cap K)$. Hence G° is a compact connected Lie group. Now we want to show the same for the group $G^\circ K$. The group $\text{Cen}_G(G^\circ)$ is normal in G and has therefore trivial intersection with K , because the action is faithful. Thus $\text{Cen}_K(G^\circ) = \{1\}$. Therefore K injects into the compact Lie group $\text{Aut}(G^\circ)$. Now both G° and $G^\circ K/G^\circ \cong K/K \cap G^\circ$ are Lie groups and thus $G^\circ K$ is also a Lie group by Lemma 2.3. Finally, $G^\circ K$ is open in G , hence G is also a Lie group. \square

3.6 Corollary *Let G be a compact group and K a closed subgroup. If $X = G/K$ is piecewise contractible, then G/K is a closed, but not necessarily connected manifold. The quotient G/N , where $N = \bigcap \{gKg^{-1} \mid g \in G\}$ is a compact Lie group that acts faithfully and transitively on X .* \square

Theorem 3.5 and Corollary 3.6 yield Theorem A in the introduction. For the sake of completeness, we restate the result in terms of transformation groups.

3.7 Corollary *Let X be a compact locally contractible space. Suppose that a compact group G acts as a transitive transformation group on X , via a continuous map*

$$G \times X \longrightarrow X.$$

Then X is a closed manifold. If the G -action is faithful, then G is a compact Lie group.

Proof. Let K denote the stabilizer of a point $x \in X$. Since G is compact, the natural continuous map $G/K \longrightarrow X$ is a homeomorphism. \square

4 A splitting result for locally compact groups

In [26], p. 547, Theorem 11, Iwasawa proved the following.

4.1 Theorem (Iwasawa's Local Splitting Theorem) *Let G be a locally compact connected group. Then G has arbitrarily small neighborhoods which are of the form NC such that N is a compact normal subgroup and C is an open n -cell which is a local Lie group commuting elementwise with N , such that $(n, c) \mapsto nc$ is a homeomorphism $N \times C \longrightarrow NC$.* \square

Iwasawa had to assume that G is a projective limit of Lie groups. However, in the process of settling Hilbert's Fifth Problem (see [29], p. 184), Yamabe showed that every locally compact group has an open subgroup which is a projective limit of Lie groups (see [29], p. 175). Thus if G is an arbitrary locally compact *connected* group we may apply Iwasawa's Local Splitting Theorem. We now show how to extend Iwasawa's Splitting Theorem to not necessarily connected locally compact groups. This result is essentially Gluškov's Theorem A in [15]. It is also proved in [20] Theorem 4.1 in a different way. We begin with two lemmas.

4.2 Lemma *Let A and B compact connected abelian groups. Suppose that we are given continuous maps $\mathbb{R}^m \xrightarrow{\varphi} B \xleftarrow{p} A$. If p is surjective, then the lifting problem*

$$\begin{array}{ccc} & & A \\ & \nearrow \tilde{\varphi} & \downarrow p \\ \mathbb{R}^m & \xrightarrow{\varphi} & B \end{array}$$

has a solution $\tilde{\varphi}$.

Proof. We dualize the diagram. The Pontrjagin duals \hat{A} and \hat{B} are discrete torsion free abelian groups and \hat{p} is injective. Moreover, $\widehat{\mathbb{R}^m} \cong \mathbb{R}^m$. The dual problem

$$\begin{array}{ccc} & & \hat{A} \\ & \nearrow \hat{\tilde{\varphi}} & \uparrow \hat{p} \\ \widehat{\mathbb{R}^m} & \xleftarrow{\hat{\varphi}} & \hat{B} \end{array}$$

clearly has a solution (for example by passing to the divisible hulls of \hat{A} and \hat{B} , which are \mathbb{Q} -vector spaces). Note that we do not have to worry about continuity, since both \hat{A} and \hat{B} are discrete groups. Now we dualize the solution $\hat{\tilde{\varphi}}$ of this problem. \square

4.3 Lemma *Let L be a simply connected Lie group and let N be a compact group. Let $\alpha : L \longrightarrow \text{Aut}(N)$ be a homomorphism, and consider the semidirect product $N \rtimes_{\alpha} L$. If L centralizes under this map the identity component N° , then there is an isomorphism*

$$\varphi : N \rtimes_{\alpha} L \xrightarrow{\cong} N \times L$$

which restricts to the identity on $N \times 1$.

Proof. The α -image of L is contained in the identity component $\text{Aut}(N)^{\circ}$, because L is connected. On the other hand, we have a natural injective map $N/\text{Cen}(N) = \text{Inn}(N) \hookrightarrow \text{Aut}(N)$. Under this map, $\text{Aut}(N)^{\circ} \cong N^{\circ}/\text{Cen}(N) \cap N^{\circ}$, see [26] p. 514, Theorem 1' or [21] Theorem 9.82 (in whose formulation the word ‘‘Lie’’ should be omitted). The subgroup of $\text{Aut}(N)^{\circ}$ that centralizes N° is therefore isomorphic to $\text{Cen}(N^{\circ})/\text{Cen}(N) \cap N^{\circ}$. Thus the L -action on N is given by a map $L \longrightarrow (\text{Cen}(N^{\circ})/\text{Cen}(N) \cap N^{\circ})^{\circ}$. The target group of this map is a quotient of the compact connected abelian group $\text{Cen}(N^{\circ})^{\circ}$. Since this group is in particular abelian, we end up with a map $L \xrightarrow{\text{ab}} L/[L, L] \xrightarrow{\varphi} (\text{Cen}(N^{\circ})/\text{Cen}(N) \cap N^{\circ})^{\circ}$. Now L is simply connected and thus $L/[L, L] \cong \mathbb{R}^m$, for some $m \geq 0$. By Lemma 4.2, there exists a lift $\tilde{\varphi} : L/[L, L] \longrightarrow \text{Cen}(N^{\circ})^{\circ}$. Now we consider the composite $\beta = j \circ \tilde{\varphi} \circ \text{ab}$,

$$L \xrightarrow{\text{ab}} L/[L, L] \xrightarrow{\tilde{\varphi}} \text{Cen}(N^{\circ})^{\circ} \xrightarrow{j} N.$$

where $j(x) = x^{-1}$. Then we have

$$\beta(g)^{-1}n\beta(g) = \alpha(g)(n)$$

for all $n \in N$ and $g \in L$. Now

$$N \times L \longrightarrow N \rtimes L, \quad \varphi(n, \ell) = (n\beta(\ell), \ell)$$

is the desired isomorphism. □

The following result, which is Theorem C in our introduction, is a global version of Gluškov’s Theorem A in [15].

4.4 Theorem *Let G be a locally compact group. Then for every identity neighborhood U there is a compact subgroup N contained in U , a simply connected Lie group L , and an open and continuous homomorphism $\varphi : N \times L \longrightarrow G$ with discrete kernel such that $\varphi(n, 1) = n$ for all $n \in N$.*

Proof. We divide the proof into several steps.

Claim 1. The result holds if G is connected. We apply Iwasawa’s Local Splitting Theorem 4.1. The fact that C is a local Lie group on an open n -cell means that there is a Lie group L , an n -cell identity neighborhood $W \subseteq L$, and a homeomorphism $\gamma : W \longrightarrow C$ for which $x, y, xy \in W$ implies $\gamma(xy) = \gamma(x)\gamma(y)$. We may assume L to be simply connected. Then γ extends to a unique homomorphism of topological groups $\gamma : L \longrightarrow G$, see [21], Corollary A2.26 and A2.27. Since C is in the centralizer of N , so is the subgroup $\alpha(L)$ generated by C . Hence the map

$$\varphi : N \times L \longrightarrow G, \quad \varphi(n, \ell) = n\gamma(\ell),$$

is a continuous homomorphism which maps $N \times W$ homeomorphically onto the identity neighborhood NC of G . Thus $\ker \varphi$ is discrete and φ is locally open and hence open. Clearly $\varphi(n, 1) = n$. The assertion follows in this special case.

Claim 2. The result holds if G/G° is compact. Then every identity neighborhood contains a compact normal subgroup P such that G/P is a Lie group, see [29] 4.6, p. 175. Let $U \subseteq G$ be an identity neighborhood. By Theorem 4.1, the identity component G° has a relatively open identity neighborhood $QC \cong Q \times C$ with a compact normal subgroup $Q \trianglelefteq G^\circ$ contained in U and an open n -cell local Lie group C . We may assume that the n -cell C contains no subgroup besides $\{1\}$. Let $\psi : Q \times L \longrightarrow G^\circ$ be the surjective homomorphism guaranteed by Claim 1 of the proof, and put

$$\gamma : L \longrightarrow G, \quad \gamma(\ell) = \psi(1, \ell).$$

Let $\mathcal{N}(G)$ be the filter basis of compact normal subgroups $P \trianglelefteq G$ such that G/P is a Lie group. Since the filter basis $\mathcal{N}(G)$ converges to 1, there is a $P \in \mathcal{N}(G)$ such that $P \subseteq U$ and $P \cap G^\circ \subseteq QC$. Since C contains no nontrivial subgroups, we conclude that $P \cap G^\circ \subseteq Q$ (because $\text{pr}_C(P \cap G^\circ)$ is a subgroup of C). Since G/P is a Lie group and G/G° is compact, G/PG° is finite. Thus PG° is open in G , and we may as well assume that $G = PG^\circ$. The group $\gamma(L)$ centralizes Q and normalizes $P \trianglelefteq G$. Therefore it normalizes the compact group $N = PQ \subseteq G$. We put

$$\alpha : L \longrightarrow \text{Aut}(N), \quad \alpha(\ell)(n) = \gamma(\ell)n\gamma(\ell)^{-1}$$

Then the semidirect product $M \rtimes_\alpha L$ has a continuous homomorphism

$$\varphi : N \rtimes_\alpha L \longrightarrow G, \quad \varphi(n, \ell) = n\gamma(\ell).$$

Its image contains P , Q and $\gamma(L)$. Therefore it maps onto $PQ\gamma(L) = PG^\circ = G$. Since L and N are σ -compact, the group $N \rtimes L$ is σ -compact. By the Open Mapping Theorem for Locally Compact Groups (see e.g. [21] p. 650), φ is open. We claim that the kernel is discrete. Let $W = \gamma^{-1}(C)$. Then $N \times W$ is an identity neighborhood of $N \rtimes_\alpha L$. Suppose that $(n, w) \in (N \times W) \cap \ker \varphi$. Then $n = \gamma(w)^{-1} = c \in C$ and $n = qp$ for some $p \in P$ and $q \in Q$. Thus $p = q^{-1}c \in QC \subseteq G^\circ$. It follows that $p \in P \cap G^\circ \subseteq Q$. Thus we may assume that $p = 1$, and this implies $n = q = c = 1$. Thus φ has a discrete kernel. Next we note that $N^\circ \subseteq G^\circ$, and thus $N^\circ \subseteq Q$. Therefore L centralizes N° . By Lemma 4.3, we have an isomorphism

$$N \times L \longrightarrow N \rtimes_\alpha L$$

which is the identity on $N \times \{1\}$. Thus we have proved Claim 2.

Claim 3. The result holds for arbitrary locally compact groups G . In such a group G , there is an open subgroup $H \subseteq G$ such that H/H° is compact. By Claim 2 we find a simply connected Lie group L , a compact subgroup N and an open homomorphism

$$N \times L \longrightarrow H \hookrightarrow G.$$

□

5 The proof of Theorem B

We now extend our results from Section 3 to locally compact groups.

5.1 Locally contractible spaces By way of comparison we remind the reader that a space X is called *locally contractible* if it satisfies the following condition.

(LC*) Every point $x \in X$ has a neighborhood V such that some open neighborhood $U \subseteq V$ of x is contractible in V .

A locally contractible space is locally arcwise connected and piecewise contractible, that is,

$$(LC^*) \implies (PC).$$

A neighborhood retract of a locally contractible space is locally contractible, see [19] Theorem 4–42 or [25] I.9. It follows that a product of two spaces is locally contractible if and only if the two factors are locally contractible. We also note that being locally contractible is a local property of a space.

In view of Theorem 4.4 the following Lemma is the main step in this extension.

5.2 Lemma (The Reduction Lemma) *Let L be a Lie group and let N be a compact group. Suppose that K is a closed subgroup of the locally compact group $G = N \times L$. If $X = G/K$ is locally contractible, then X is a manifold. If N acts faithfully on X , then N is a compact Lie group and hence G is a Lie group.*

Proof. Since N is compact and normal in G , the group $NK \subseteq G$ is closed, see [18] II 4.4. Because N is a direct factor in G , the group NK splits as $NK = N \times H$, and $H \subseteq L$ is a closed Lie subgroup. The natural map

$$\begin{array}{ccc} N \times H & \longrightarrow & N \times L \\ & & \downarrow \\ & & (N \times L)/(N \times H) \end{array}$$

is a locally trivial principal bundle because this is true for the map $L \longrightarrow L/H$, see [37], Theorem 3.58. It follows that the associated bundle

$$\begin{array}{ccc} (N \times H)/K & \longrightarrow & (N \times L)/K \\ & & \downarrow \\ & & (N \times L)/(N \times H) \end{array}$$

is also locally trivial. Since $(N \times L)/K$ is locally contractible, the same is true for the fiber $(N \times H)/K$ by 5.1. Now the compact group N acts transitively on the fiber

$$F = (N \times H)/K = NK/K \cong N/N \cap K.$$

By Theorem 3.5, the homogeneous space F is a closed manifold. If N acts faithfully on G/K , then N acts faithfully on F , because a subgroup $P \subseteq N \cap K$ which is normal in N is also normal in $N \times L$. Hence N is a compact Lie group in this case. \square

5.3 Corollary *Let G be a locally compact group and let $K \subseteq G$ be a closed subgroup. If $X = G/K$ is locally contractible, then X is a manifold.*

Proof. Let $\varphi : N \times L \longrightarrow G$ be as in Theorem 4.4. Then $H = \varphi(N \times L)$ is an open subgroup of G . Therefore H has an open orbit $Y = HK/K \subseteq X$. By Lemma 5.2, this open set Y is a manifold. It follows from the homogeneity of X that X itself is a manifold. \square

The following consequence is immediate. This is Theorem B in the introduction.

5.4 Theorem *Let G be a locally compact group and let $K \subseteq G$ be a closed subgroup. If $X = G/K$ is locally contractible and connected and if G acts faithfully on X , then G is a Lie group.*

Proof. We argue similarly as in the proof of Corollary 5.3 and we consider the open subgroup $H = \varphi(N \times L)$. The H -orbit of the coset $gK \in G/K$ is the open set HgK/K . Because X is connected, this implies that H acts transitively on X . In particular, $N \subseteq H$ acts faithfully on $H/H \cap K \cong G/K$. By Lemma 5.2, H is a Lie group. Since $H \subseteq G$ is open, G is also a Lie group. \square

Again, we restate this result in terms of transformation groups.

5.5 Corollary *Let G be a locally compact and σ -compact group. If X is a locally contractible, locally compact and connected space and if $G \times X \longrightarrow X$ is a continuous faithful action, then G is a Lie group and X is a manifold.*

Proof. Let $x \in X$ be a point and let $K \subseteq G$ denote the stabilizer of x . Since G is σ -compact, the natural continuous map $G/K \longrightarrow X$ is a homeomorphism, see for example [35] 10.10. \square

6 A historical review

In his influential 1974 paper [36] Szenthe stated the following result on locally compact groups:

Theorem 4 [36]. Let a σ -compact group G with compact G/G° be an effective and transitive topological transformation group of a locally compact and locally contractible space X . The G is a Lie group and X is homeomorphic to a coset space of G .

Szenthe's statement provided a result which was needed and applied in various areas, notably in geometry, see for example [1, 4, 5, 34, 16, 17, 27]. The proof of Theorem 4 was based, among other things, on the following statement on compact groups.

Lemma 6 [36]. Let G be a compact group, $H \subseteq G$ a closed subgroup, $\chi : G \longrightarrow G/H$ the canonical projection, $A \subseteq G$ a closed invariant subgroup and $A' = \chi(A)$. If A' is contractible over G/H , then $A \subseteq H$.

However, in 2011, Sergey Antonyan [3] found the following simple counterexample to Szenthe's Lemma 6:

Example. Let $G = \mathbb{S}^1 \subseteq \mathbb{C}^*$, let $H = \{1\}$ and put $A = \{\pm 1\}$. Then $A' = A$ is contractible in G , but is not contained in H .

It was also noted the mid-nineties by Salzmänn and his school [34] that Szenthe's method of approximating a locally compact group G by Lie groups forces G to be metric, that is, first countable, see [6]. Thus even if a substitute method for Szenthe's Lemma 6 for compact groups could be obtained by some correct argument, the ensuing version of Theorem 4 could only be valid for first countable locally compact groups.

References

- [1] S. M. Ageev and D. Repovš, *Mat. Sb.* **201** (2010), no. 2, 3–28; translation in *Sb. Math.* **201** (2010), no. 1-2, 159–182. MR2656322 (2011h:54050)
- [2] S. A. Antonyan, Equivariant extension properties of coset spaces of locally compact groups and approximate slices, *Topology Appl.* **159** (2012), no. 9, 2235–2247. MR2921813
- [3] S. A. Antonyan, Characterizing maximal compact subgroups, *Arch. Math. (Basel)* **98** (2012), no. 6, 555–560. MR2935661
- [4] V. N. Berestovskii, *Izv. Vyssh. Uchebn. Zaved. Mat.* **2004**, no. 11, 3–22; translation in *Russian Math. (Iz. VUZ)* **48** (2004), no. 11, 1–19 (2005). MR2179443 (2006i:53055)
- [5] V. N. Berestovskii, D. M. Halverson and D. Repovš, Locally G -homogeneous Busemann G -spaces, *Differential Geom. Appl.* **29** (2011), no. 3, 299–318. MR2795840 (2012g:53153)
- [6] H. Bickel, Lie-projective groups, *J. Lie Theory* **5** (1995), no. 1, 15–24. MR1362007 (96k:22009)
- [7] G. E. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972. MR0413144 (54 #1265)
- [8] J. D. Dixon and B. Mortimer, *Permutation groups*, Graduate Texts in Mathematics, 163, Springer, New York, 1996. MR1409812 (98m:20003)
- [9] J. Dugundji, *Topology*, Allyn & Bacon, Boston, MA, 1978. MR0478089 (57 #17581)
- [10] S. Eilenberg and N. Steenrod, *Foundations of algebraic topology*, Princeton Univ. Press, Princeton, New Jersey, 1952. MR0050886 (14,398b)
- [11] R. Engelking, *General topology*, translated from the Polish by the author,
- [12] A. A. George Michael, A counterexample in the dimension theory of homogeneous spaces of locally compact groups, *J. Lie Theory* **18** (2008), no. 4, 915–917. MR2523143 (2010b:22010)
- [13] A. A. George Michael, On three classical results on compact groups, *Scientiae Mathematicae Japonica Online*, e-2011, 271–273.

- [14] A. A. George Michael, submitted.
- [15] V. M. Gluškov, Structure of locally bicomact groups and Hilbert's fifth problem, *Uspehi Mat. Nauk (N.S.)* **12** (1957), no. 2 (74), 3–41. MR0101892 (21 #698)
Translated as:
V. M. Gluškov, The structure of locally compact groups and Hilbert's fifth problem, *Amer. Math. Soc. Transl. (2)* **15** (1960), 55–93. MR0114872 (22 #5690)
- [16] T. Grundhöfer, N. Knarr and L. Kramer, Flag-homogeneous compact connected polygons, *Geom. Dedicata* **55** (1995), no. 1, 95–114. MR1326742 (96a:51009)
- [17] T. Grundhöfer, N. Knarr and L. Kramer, Flag-homogeneous compact connected polygons. II, *Geom. Dedicata* **83** (2000), no. 1-3, 1–29. MR1800008 (2001m:51022)
- [18] E. Hewitt and K. A. Ross, *Abstract harmonic analysis. Vol. I*, second edition,
- [19] J. G. Hocking and G. S. Young, *Topology*, Addison-Wesley Publishing Co., Inc., Reading, MA, 1961. MR0125557 (23 #A2857)
- [20] K. H. Hofmann and S. A. Morris, Transitive actions of compact groups and topological dimension, *J. Algebra* **234** (2000), no. 2, 454–479. MR1801101 (2002a:22006)
- [21] K. H. Hofmann and S. A. Morris, *The structure of compact groups*, de Gruyter Studies in Mathematics, 25, de Gruyter, Berlin, 1998. MR1646190 (99k:22001)
- [22] K. H. Hofmann and S. A. Morris, *The Lie theory of connected pro-Lie groups*, EMS Tracts in Mathematics, 2, European Mathematical Society (EMS), Zürich, 2007. MR2337107 (2008h:22001)
- [23] K. H. Hofmann and P. S. Mostert, *Elements of compact semigroups*, Charles E. Merril Books, Inc., Columbus, OH, 1966. MR0209387 (35 #285)
- [24] K. H. Hofmann and K.-H. Neeb, Pro-Lie groups which are infinite-dimensional Lie groups, *Math. Proc. Cambridge Philos. Soc.* **146** (2009), no. 2, 351–378. MR2475971 (2009k:22028)
- [25] S. Hu, *Theory of retracts*, Wayne State Univ. Press, Detroit, 1965. MR0181977 (31 #6202)
- [26] K. Iwasawa, On some types of topological groups, *Ann. of Math. (2)* **50** (1949), 507–558. MR0029911 (10,679a)
- [27] L. Kramer, Homogeneous spaces, Tits buildings, and isoparametric hypersurfaces, *Mem. Amer. Math. Soc.* **158** (2002), no. 752, xvi+114 pp. MR1904708 (2003m:53078)
- [28] B. Madison, Semigroups on coset spaces, *Duke Math. J.* **36** (1969), 61–63. MR0237702 (38 #5983)
- [29] D. Montgomery and L. Zippin, *Topological transformation groups*, Interscience Publishers, New York, 1955. MR0073104 (17,383b)

- [30] Math Reviews MR0578648 (58 #28253) Albu, A. C., Sur un lemma de J. Szenthe [36].
- [31] Math Reviews MR0360921 (50 #13368) K. H. Hofmann on: [36]
- [32] A. L. Onishchik, *Topology of transitive transformation groups*, Johann Ambrosius Barth, Leipzig, 1994. MR1266842 (95e:57058)
- [33] L. S. Pontrjagin, *Topologische Gruppen. I*, B. G. Teubner Verlagsgesellschaft, Leipzig, 1957. MR0086265 (19,152e)
- [34] H. Salzmann et al., *Compact projective planes*, de Gruyter Expositions in Mathematics, 21, de Gruyter, Berlin, 1995. MR1384300 (97b:51009)
- [35] M. Stroppel, *Locally compact groups*, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2006. MR2226087 (2007d:22001)
- [36] J. Szenthe, On the topological characterization of transitive Lie group actions, Acta Sci. Math. (Szeged) **36** (1974), 323–344. MR0360921 (50 #13368)
- [37] F. W. Warner, *Foundations of differentiable manifolds and Lie groups*, corrected reprint of the 1971 edition, Graduate Texts in Mathematics, 94, Springer, New York, 1983. MR0722297 (84k:58001)
- [38] H. Yamabe, On the conjecture of Iwasawa and Gleason, Ann. of Math. (2) **58** (1953), 48–54. MR0054613 (14,948e)

Karl H. Hofmann
 Technische Universität Darmstadt
 Schlossgartenstraße 7
 64289 Darmstadt, Germany
 hofmann@mathematik.tu-darmstadt.de

Linus Kramer
 Mathematisches Institut
 Einsteinstr. 62
 48149 Münster, Germany
 linus.kramer@uni-muenster.de